

Dynamic oxygen mapping in tumors by ^{19}F MRI

Steven Baete

Supervisor(s): Yves De Deene

Tumour hypoxia is well known to reduce cancer treatment efficacy. Hypoxic tumour cells, which have decreased oxygen levels ($p\text{O}_2$), are more resistant to radiotherapy and chemotherapy. Identification of hypoxic tumour tissue may improve cancer treatment. The oxygen levels in tissue can be imaged in a non-invasive way by ^{19}F MRI oximetry. That is, one of the parameters which can be measured quantitatively with magnetic resonance imaging (MRI), the spin-lattice relaxation rate R_1 ($R_1 = 1 / T_1$) of perfluorocarbons, is sensitive to changes in oxygen tension $p\text{O}_2$ [1]. This technique has been used in animal studies to dynamically image tumour oxygenation (e.g. [1]). However, validation of $p\text{O}_2$ -measurements in vivo is difficult. In this study a reproducible phantom, simulating well perfused oxygen consuming tissue, is presented. Furthermore, computer simulations are used to study the impact of perfluorocarbon distribution and concentration in tissue on ^{19}F MRI oximetry.

I. PHANTOM

A phantom (figure 1)[2] was used to simulate and image acute and perfusion-related hypoxia spatially and temporally. This phantom can be employed to simulate oxygen consumption by somatic cells in vivo and for validating computational biophysical models of hypoxia, as measured with ^{19}F MRI oximetry.

II. SIMULATION

A Krogh tissue model is implemented. This model, in which a tissue cylinder surrounds a capillary, incorporates the presence of perfluorocarbons in blood and tissue. Mean oxygen tension $p\text{O}_2$ in the simulation volume is compared with $p\text{O}_2$ as it would be measured by ^{19}F




Figure 1. Schematic of a phantom for the validation of ^{19}F MRI-oximetry methods. The phantom consists of a hemodialysis filter placed in an MR-scanner and an automated syringe pump. The outer compartment of the filter is filled with a gelatin matrix containing viable yeast cells simulating oxygen consuming tissue. Also perfluorocarbon emulsions can be added to the gelatin matrix.

MRI oximetry, $p\text{O}_2^{\text{FNMR}}$.

Simulation results indicate that the agreement between measured $p\text{O}_2^{\text{FNMR}}$ and actual $p\text{O}_2$ is influenced by vascular density, perfluorocarbon distribution and perfluorocarbon concentration. Only homogeneous perfluorocarbon distribution in tissue guarantees small deviations of $p\text{O}_2^{\text{FNMR}}$ from $p\text{O}_2$. Hence, the perfluorocarbon distribution and concentration have a serious impact on the accuracy of ^{19}F MRI oximetry.

ACKNOWLEDGMENTS

This research is supported by grants #01J13507 and #01D21208 from the BOF-Ghent University. The author is a PhD Fellow of the Research Foundation - Flanders (FWO).

REFERENCES

- [1] V. D. Kodibagkar, X. Wang, and R. P. Mason, "Physical principles of quantitative nuclear magnetic resonance oximetry," *Front. Biosci.*, vol. 13, pp. 1371–1384, 2008.
- [2] S. Baete, J. Vandecasteele, L. Colman, W. De Neve, and Y. De Deene, "An oxygen-consuming phantom simulating perfused tissue to explore oxygen dynamics and ^{19}F mri oximetry," *Mang. Reson. Mater. Phys.*, vol. 23, pp. 217–226, 2010.

S. Baete is with the Dept. for Radiation Oncology and Experimental Cancer Research, Ghent University (UGent), Gent, Belgium. E-mail: Steven.Baete@UGent.be .